Brain Research, 461 (1988) 407–409
Elsevier

Author Index

Aldskogius, H., Wiesenfeld-Hallin, Z. and Kristensson, K., Selective neuronal destruction by *Ricinus communis* agglutinin I and its use for quantitative determination of sciatic nerve dorsal root ganglion cell numbers, 215

Baranyi, A., See Szente, M.B., 64
Beesley, P.W., see Hill, I.E., 27
Ben-Ari, Y., see Tremblay, E., 393
Bigelow, J.C., see Kristensen, E.W., 44
Billings-Gagliardi, S., see Wolf, M.K., 257
Bing, G., see Kordower, J.H., 221
Björklund, A., see Nothias, F., 349
Blessing, W.W., see Li, Y.W., 75
Brödemann, R., see Reymann, K.G., 388
Byers, M.R. and Costello, R.J., Membrane heterogeneity in sensory receptors of rat vibrissae: Merkel receptors differ from other nerve endings, 397
Byers, M.R., see Taylor, P.E., 371
Byrne, M.C., Farooq, M., Sbaschnig-Agler, M., Norton, W.T. and Ledeen, R.W., Ganglioside content of astroglia and neurons isolated from maturing rat brain: consideration of the source of astroglial gangliosides, 87
Byrne, M.C., see Sbaschnig-Agler, M., 98

Cao, C.W., see Shigenaga, Y., 238
Carlsson, A., see Maysinger, D., 355
Chan, S.O. and Jen, L.S., Enlargement of uncrossed retinal projection in the albino rat: additive effects of neonatal eye removal and thalamectomy, 163
Charriaut-Marlangue, C., see Tremblay, E., 393
Coggeshall, R.E., see Hulsebosch, C.E., 322
Consolo, S., see Tacconi, M.T., 194
Costall, B., see Gibb, W.R.G., 361
Costello, R.J., see Byers, M.R., 397
Crowley, W.R., Sex differences in the response of hypothalamic luteinizing hormone-releasing hormone and catecholamine systems to ovarian hormones and naloxone: implications for sexual differentiation of luteinizing hormone secretion in rats, 314
Cuello, A.C., see Maysinger, D., 355

Davrainville, J.L., see Math, F., 384
Dickenson, A.H., see Sullivan, A.F., 182
Doe, K., see Shigenaga, Y., 143
Domeney, A.M., see Gibb, W.R.G., 361
Dragnic, Y., see Villanueva, S., 377
Dreyfus, H., see Sbaschnig-Agler, M., 98
Dubovy, P., Dipeptidylpeptidase IV activity in Meissner corpuscles of rhesus monkey and its possible function, 186

Erzurumlu, R.S., Jhaveri, S. and Schneider, G.E., Distribution of morphologically different retinal axon terminals in the hamster dorsal lateral geniculate nucleus, 175

Farooq, M., see Byrne, M.C., 87
Farooq, M., see Sbaschnig-Agler, M., 98
Fiandaca, M.S., see Kordower, J.H., 221
Fisone, G., see Tacconi, M.T., 194
Frenz, P., see Villanueva, S., 377

Garofalo, L., see Maysinger, D., 355
Gash, D.M., see Kordower, J.H., 221
Geffen, L.B., see Li, Y.W., 75
George, S.R., see Haas, D.A., 230
Gibb, W.R.G., Costall, B., Domeney, A.M., Kelly, M.E. and Naylor, R.J., The histological effects of intracerebral injection or infusion of MPTP (1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine) and MPP^+ (1-methyl-4-phenylpyridinium) in rat and mouse, 361

Haas, D.A. and George, S.R., Single or repeated mild stress increases synthesis and release of hypothalamic corticotropin-releasing factor, 230
Halliday, G.M., see Li, Y.W., 75
Hattori, Y., see Moriwaki, A., 190
Hawkes, R.B., see Hill, I.E., 27
Herranz, A.S., see Herreras, O., 303
Herrera-Marschitz, M., see Maysinger, D., 355
Herreras, O., Solís, J.M., Herranz, A.S., Martín del Río, R. and Lerma, J., Sensory modulation of hippocampal transmission. II. Evidence for a cholinergic locus of inhibition in the Schaffer–CA1 synapse, 303
Herreras, O., Solís, J.M., Muñoz, M.D., Martín del Río, R. and Lerma, J., Sensory modulation of hippocampal transmission. I. Opposite effects on CA1 and dentate gyrus synapses, 290
Hill, I.E., Selkirk, C.P., Hawkes, R.B. and Beesley, P.W., Characterization of novel glycoprotein components of synaptic membranes and postsynaptic densities, gp65 and gp55, with a monoclonal antibody, 27
Hiner, B.C., Mauk, M.D., Peroutka, S.J. and Kocsis, J.D., Buspirone, 8-OH-DPAT and ipsapirone: effects on hippocampal cerebellar and sciatic fiber excitability, 1
Hirai, K. and Katayama, Y., Late slow outward currents following hyperpolarizing commands are associated with intracellular Ca^{2+} increase in guinea-pig submucous neurons: investigations with fura-2, 367
Hökfelt, T., see Millhorn, D.E., 169
Hokoc, J.N. and Mariani, A.P., Synapses from bipolar cells onto dopaminergic amacrine cells in cat and rabbit retinas, 17
Hong, J.S., see Zhang, W.Q., 335
Hori, Y., see Moriwaki, A., 190
Hubbard, J.I., see Sibbald, J.R., 205
Hudson, P.M., see Zhang, W.Q., 335
Hulsebosch, C.E. and Coggeshall, R.E., Intraspinal sprouting after administration of nerve growth factor antibodies to neonatal rats, 322

Isacson, O., see Nothias, F., 349

Jackson, I.M.D., see Van den Bergh, P., 53
Jen, L.S., see Chan, S.O., 163
Jhaveri, S., see Erzurumlu, R.S., 175

Joh, T.H., see Li, Y.W., 75

Kai, Y., Oomura, Y. and Shimizu, N., Responses of rat lateral hypothalamic neurons to periaqueductal gray stimulation and nociceptive stimuli, 107
Kase, H., see Reymann, K.G., 388
Katayama, Y., see Hirai, K., 367
Kavaliers, M., see Wiebe, J.P., 150
Kawai, Y., see Kohno, J., 158
Kawasaki, K., Matsushita, A., Satoh, M. and Takagi, H., The lateral vestibular nucleus is a site for the depressant action of benzodiazepines on the crossed extensor reflex, 282
Kelly, M.E., see Gibb, W.R.G., 361
Khayari, A., see Math, F., 384
Klockgether, T., Turski, L., Schwarz, M., Sontag, K.-H. and Lehmann, J., Paradoxical convulsant action of a novel non-competitive *N*-methyl-D-aspartate (NMDA) antagonist, tiletamine, 343
Kocsis, J.D., see Hiner, B.C., 1
Kohno, J., Shinoda, K., Kawai, Y., Peng, Y., Ono, K. and Shiotani, Y., Direct adrenergic inputs to sacral autonomic neurons: using a double-immunostaining method at the light and electron microscopic levels, 158
Kordower, J.H., Sladek Jr., J.R., Fiandaca, M.S., Bing, G. and Gash, D.M., Tyrosine hydroxylase-immunoreactive somata within the primate subfornical organ: species specificity, 221
Kristensen, E.W., Bigelow, J.C. and Wightman, R.M., Time resolved dopamine overflow synaptosomes and chopped striatal tissue with rapid superfusion, 44
Kristensson, K., see Aldskogius, H., 215

Lechan, R.M., see Van den Bergh, P., 53
Ledeen, R.W., see Byrne, M.C., 87
Ledeen, R.W., see Sbaschnig-Agler, M., 98
Lehmann, J., see Klockgether, T., 343
Lerma, J., see Herreras, O., 290
Lerma, J., see Herreras, O., 303
Li, Y.W., Halliday, G.M., Joh, T.H., Geffen, L.B. and Blessing, W.W., Tyrosine hydroxylase-containing neurons in the supraoptic and paraventricular nuclei of the adult human, 75
Low, P.A., see Zochodne, D.W., 10
Lupa, M.T., Effects of an inhibitor of the synaptic vesicle acetylcholine transport system on quantal neurotransmitter release: an electrophysiological study, 118

Mariani, A.P., see Hokoc, J.N., 17
Martín del Río, R., see Herreras, O., 290
Martín del Río, R., see Herreras, O., 303
Math, F., Khayari, A. and Davrainville, J.L., Influence of GABA, baclofen and bicuculline on the extracellular K^+ activity in the olfactory bulb of the developing rat, 384
Matsushita, A., see Kawasaki, K., 282
Matthies, H., see Reymann, K.G., 388
Mauk, M.D., see Hiner, B.C., 1
Maysinger, D., Herrera-Marschitz, M., Carlsson, A., Garofalo, L., Cuello, A.C. and Ungerstedt, U., Striatal and cortical acetylcholine release in vivo in rats with unilateral decortication: effects of treatment with monosialoganglioside GM1, 355
McGeer, E.G., see Mizukawa, K., 274
McGeer, P.L., see Mizukawa, K., 274
Millhorn, D.E., Hökfelt, T., Seroogy, K. and Verhofstad, A.A.J., Extent of colocalization of serotonin and GABA in neurons of the ventral medulla oblongata in rat, 169
Mitsuhiro, Y., see Shigenaga, Y., 143
Mitsuhiro, Y., see Shigenaga, Y., 238
Mizukawa, K., Vincent, S.R., McGeer, P.L. and McGeer, E.G., Reduced nicotinamide adenine dinucleotide phosphate (NADPH)-diaphorase-positive neurons in cat cerebral white matter, 274
Moriwaki, A., Hattori, Y., Yasuhara, H. and Hori, Y., Lateral difference in responsiveness of norepinephrine-sensitive cyclic AMP-generating systems of rat cerebral cortex with iron-induced epileptic activity, 190
Muñoz, M.D., see Herreras, O., 290

Nanry, K.P., see Zhang, W.Q., 335
Naylor, R.J., see Gibb, W.R.G., 361
Nicholson, C., see Rice, M.E., 328
Nitsch, C. and Riesenberg, R., Immunocytochemical demonstration of GABAergic synaptic connections in rat substantia nigra after different lesions of the striatonigral projection, 127
Norton, W.T., see Byrne, M.C., 87
Norton, W.T., see Sbaschnig-Agler, M., 98
Nothias, F., Wictorin, K., Isacson, O., Björklund, A. and Peschanski, M., Morphological alteration of thalamic afferents in the excitoxically lesioned striatum, 349

Oka, H., see Yamamoto, T., 199
Ono, K., see Kohno, J., 158
Oomura, Y., see Kai, Y., 107
Orrego, F., see Villanueva, S., 377
Otani, K., see Shigenaga, Y., 238

Peng, Y., see Kohno, J., 158
Peroutka, S.J., see Hiner, B.C., 1
Peschanski, M., see Nothias, F., 349

Redd, P.E., see Taylor, P.E., 371
Represa, A., see Tremblay, E., 393
Reymann, K.G., Brödemann, R., Kase, H. and Matthies, H., Inhibitors of calmodulin and protein kinase C block different phases of hippocampal long-term potentiation, 388
Rice, M.E. and Nicholson, C., Behavior of extracellular K^+ and pH in skate (*Raja erinacea*) cerebellum, 328
Riesenberg, R., see Nitsch, C., 127
Roisin, M.P., see Tremblay, E., 393

Saito, H., see Tanaka, J., 403
Samejima, A., see Yamamoto, T., 199
Satoh, M., see Kawasaki, K., 282
Sbaschnig-Agler, M., Dreyfus, H., Norton, W.T., Sensenbrenner, M., Farooq, M., Byrne, M.C. and Ledeen, R.W., Gangliosides of cultured astroglia, 98
Sbaschnig-Agler, M., see Byrne, M.C., 87
Schneider, G.E., see Erzurumlu, R.S., 175
Schwarz, M., see Klockgether, T., 343
Scott, J.J.A., Responses of cat muscle spindle afferents after short periods of denervation and reinnervation, 381
Selkirk, C.P., see Hill, I.E., 27
Sensenbrenner, M., see Sbaschnig-Agler, M., 98
Seroogy, K., see Millhorn, D.E., 169
Seto, K., see Tanaka, J., 403
Shigenaga, Y., Yoshida, A., Mitsuhiro, Y., Tsuru, K. and Doe, K., Morphological and functional properties of trigeminal nucleus oralis neurons projecting to the trigeminal mo-

tor nucleus of the cat, 143
Shigenaga, Y., Yoshida, A., Tsuru, K., Mitsuhiro, Y., Otani, K. and Cao, C.Q., Physiological and morphological characteristics of cat masticatory motoneurons — intracellular injection of HRP, 238
Shimizu, N., see Kai, Y., 107
Shinoda, K., see Kohno, J., 158
Shiotani, Y., see Kohno, J., 158
Sibbald, J.R., Hubbard, J.I. and Sirett, N.E., Responses from osmosensitive neurons of the rat subfornical organ in vitro, 205
Sirett, N.E., see Sibbald, J.R., 205
Sladek Jr., J.R., see Kordower, J.H., 221
Solís, J.M., see Herreras, O., 290
Solís, J.M., see Herreras, O., 303
Sontag, K.-H., see Klockgether, T., 343
Stachowiak, M.K., see Zhang, W..Q., 335
Sullivan, A.F. and Dickenson, A.H., Electrophysiological studies on the spinal effects of dermorphin, an endogenous μ-opioid agonist, 182
Szente, M.B., Baranyi, A. and Woody, C.D., Intracellular injection of apamin reduces a slow potassium current mediating afterhyperpolarizations and IPSPs in neocortical neurons of cats, 64

Tacconi, M.T., Fisone, G. and Consolo, S., Modulation of phospholipid methylation in rat striatum by the corticostriatal pathway, 194
Takagi, H., see Kawasaki, K., 282
Tanaka, J., Saito, H. and Seto, K., The median preoptic nucleus participates in the control of paraventricular vasopressin neurons by the subfornical organ in the rat, 403
Taylor, P.E., Byers, M.R. and Redd, P.E., Sprouting of CGRP nerve fibers in response to dentin injury in rat molars, 371
Tilson, H.A., see Zhang, W.Q., 335
Tremblay, E., Roisin, M.P., Represa, A., Charriaut-Marlangue, C. and Ben-Ari, Y., Transient increased density of NMDA binding sites in the developing rat hippocampus, 393
Tsuru, K., see Shigenaga, Y., 143
Tsuru, K., see Shigenaga, Y., 238
Turski, L., see Klockgether, T., 343

Ungerstedt, U., see Maysinger, D., 355

Van den Berg, P., Wu, P., Jackson, I.M.D. and Lechan, R.M., Neurons containing a N-terminal sequence of the TRH-prohormone (preproTRH$_{53-74}$) are present in a unique location of the midbrain periaqueductal gray of the rat, 53
Verhofstad, A.A.J., see Millhorn, D.E., 169
Villanueva, S., Frenz, P., Dragnic, Y. and Orrego, F., Veratridine-induced release of endogenous glutamate from rat brain cortex slices: a reappraisal of the role of calcium, 377
Vincent, S.R., see Mizukawa, K., 274

Ward, K.K., see Zochodne, D.W., 10
Wictorin, K., see Nothias, F., 349
Wiebe, J.P. and Kavaliers, M., Analgesic effects of the putative FSH-suppressing gonadal steroid, 3α-hydroxy-4-pregnen-20-one: possible modes of action, 150
Wiesenfeld-Hallin, Z., see Aldskogius, H., 215
Wightman, R.M., see Kristensen, E.W., 44
Wolf, M.K. and Billings-Gagliardi, S., Quaking*shiverer double mutant mice: morphological phenotypes support possible dual actions of the shiverer locus, 257
Woody, C.D., see Szente, M.B., 64
Wu, P., see Van den Bergh, P., 53

Yamamoto, T., Samejima, A. and Oka, H., Short latency activation of local circuit neurons in the cat somatosensory cortex, 199
Yasuhara, H., see Moriwaka, A., 190
Yoshida, A., see Shigenaga, Y., 143
Yoshida, A., see Shigenaga, Y., 238

Zhang, W.Q., Tilson, H.A., Nanry, K.P., Hudson, P.M., Hong, J.S. and Stachowiak, M.K., Increased dopamine release from striata of rats after unilateral nigrostriatal bundle damage, 335
Zochodne, D.W., Ward, K.K. and Low, P.A., Guanethidine adrenergic neuropathy: an animal model of selective autonomic neuropathy, 10